



eISSN 2330-7250

JOURNAL HOME PAGE AT [WWW.VEDICJOURNALS.COM](http://WWW.VEDICJOURNALS.COM)

VRI Press

## BIOLOGICAL MEDICINAL CHEMISTRY

## SHORT COMMUNICATION

DOI: <http://dx.doi.org/10.14259/bmc.v3i1.157>

## Cardiovascular Diseases Correlation with Antioxidants in Hilla City, Iraq

MUFEED J EWADH\*, MONEM M ALSHOK, MENA Y ABID

College of Medicine, University of Babylon, Hilla, Iraq.

Article Info: Received: June 26th, 2014; Accepted: December 10th, 2014

## ABSTRACT

Cardiovascular disease (CVD) covers any disease of the circulatory system, namely the heart (cardio) or blood vessels (vascular), includes heart attack, angina and peripheral vascular heart disease, however antioxidants are vital substances that possess the ability to protect the body from damage caused by free radicals induce oxidative stress. This study aimed to find out the possible mechanisms of correlation between CVD and antioxidants. The results showed that total protein (TP), total antioxidant status (TAS), albumin levels, catalase (CAT), superoxide dismutase (SOD) have significance decreased in patients compared to control the group.

**Keywords:** Cardiovascular diseases, Antioxidants, Catalase, Total protein, Serum albumine

## Introduction

Free radicals are very reactive chemical species with an unpaired electron. Because of their reactivity lipids, proteins and DNA can be damaged by the radical's action. In consequence, they are responsible for many diseases, such as cancer, cardiovascular disorder, atherosclerosis, asthma, arthritis, neurodegenerative disorders: Alzheimer's, Parkinson's diseases and dementia [1]. As a first level of defense, the AOS includes enzymes, mostly intracellular located, such as several forms of superoxide dismutase (SOD), catalase (CAT) and a group of macromolecules including albumin [2]. The chemistry of antioxidants involves the mechanism of action of antioxidant that two mechanisms of action have been proposed for antioxidants. The first is a chain-breaking mechanism by which the primary antioxidants donate electrons to the free radicals present in the system, example lipid radicals. The second mechanism involves removal of ROS (reactive oxygen species) and RNS (reactive nitrogen species) initiator by quenching chain initiator catalyst [3]. On this basis, there are five known antioxidant mechanisms which describe antioxidant reactions [4, 5].

\*Corresponding Author

Prof. Mufeed J Ewadh

College of Medicine, University of Babylon, Hilla, Iraq.

Email: [mewadh@yahoo.com](mailto:mewadh@yahoo.com)

## Materials and Methods

1. Patients with AMI (n= 60), Control (n= 40)
2. All chemicals used in this study with highly purified and no further purification done.

## Collection of specimens:

Venous blood samples of 10 milliliters were drawn from patients of AMI within 12 hours of the heart attacks by using disposable syringes (10mL) in sitting position (Ethical committee certificate No.5/2014), 1.8 mL of fresh blood drawn in a plane tube contains (0.2mL) trisodium citrate solution 3.8% as anticoagulant for measuring fibrinogen, the remaining of the blood sample pushed slowly into plain disposable tubes. The blood was allowed to clot at 37° C for 15 minutes, and then centrifuged at 3000 rpm for 10 minutes then the sera were obtained and stored at -20° C until analysis of TP, TAS, SOD, CAT, Albumin level.

## Determination of Total Protein

Serum TP was measured using kit supplied by Spinreact (Spain) [6]

## Determination of TAS

Serum TAS was measured using kit supplied by Bio cat. (Germany) [7]



**Determination of SOD**

(SOD) activity measured using the method as described previously [7]

**Determination of CAT**

(CAT) activity measured using the method as described by Cohen et al., [8]

**Determination of Serum albumin :**

Serum albumin was measured by colorimetric assay using kit supplied by Manufacturer Biolabo (France) [9]

concentration is the result of the fall in albumin, that A: G ratio change due to either reduction of albumin or elevation of globulins, however, decreased levels (hypoproteinemia) are found in Renal diseases, malnutrition, albuminuria, and terminal liver failure. Serum albumin is a negative acute-phase reactant that is synthesized by the liver, in inflammatory states, the activity of macrophages and other immune system cells is enhanced, leading to the production of cytokines, which switch protein synthesis in the liver from serum albumin to other acute-phase proteins. The albumin-MI association was modified by hypertension, in those hypertensive subjects showed a steeper relation of albumin to MI risk. There was evidence for an increased risk of all-cause mortality with lower albumin levels among women (R1). Albumin at concentrations less than physiological can inhibit copper-stimulated peroxidation and hemolysis. It also inhibits the production of free hydroxyl radicals from systems containing copper ions and  $H_2O_2$  and can scavenge peroxy radicals, and its inhibits copper

## Results and Discussion

The results of this study (Tables 1-5) showed a significant decrease in total protein levels in Patients group compared with the control group ( $P<0.001$ ). A decrease in total protein

**Table 1: Serum levels of Serum Total Protein (gm/dl) of patients and control**

| Samples  | Number | Total Protein (gm/dl)<br>Mean $\pm$ SD | P value   |
|----------|--------|----------------------------------------|-----------|
| Control  | 40     | 7.7938 $\pm$ 0.33092                   | $P<0.001$ |
| Patients | 60     | 5.6752 $\pm$ 0.49884                   |           |

**Table 2: Serum levels of Serum Albumin (gm/dl) with AMI patients and controls**

| Samples  | Number | Serum Albumin (gm/dl)<br>Mean $\pm$ SD | P value   |
|----------|--------|----------------------------------------|-----------|
| Control  | 40     | 4.5420 $\pm$ 0.1283                    | $P<0.001$ |
| Patients | 60     | 3.4065 $\pm$ 0.18431                   |           |

**Table 3: Serum levels of SOD (U/L) with AMI patients and controls**

| Samples  | Number | SOD (U/mL)<br>Mean $\pm$ SD | P value   |
|----------|--------|-----------------------------|-----------|
| Control  | 40     | 23.5925 $\pm$ 2.12885       | $P<0.001$ |
| Patients | 60     | 16.8517 $\pm$ 0.78167       |           |

**Table 4: Serum levels of TAC (mmol/L) with AMI patients and controls**

| Samples  | Number | TAC<br>Mean $\pm$ SD | P value   |
|----------|--------|----------------------|-----------|
| Control  | 40     | 3.5450 $\pm$ 0.75037 | $P<0.001$ |
| Patients | 60     | 1.3520 $\pm$ 0.33081 |           |

**Table 5: Serum levels of CAT (U/L) with AMI patients and controls**

| Samples  | Number | TAC<br>Mean $\pm$ SD | P value   |
|----------|--------|----------------------|-----------|
| Control  | 40     | 3.5450 $\pm$ 0.75037 | $P<0.001$ |
| Patients | 60     | 1.3520 $\pm$ 0.33081 |           |

dependent lipid peroxidation systems. LDL oxidation is one of the early steps in the atherosclerotic process that serum albumin may inhibit endothelial apoptosis [10]. The result is in agreement with results of [11,12] (Table 1, 2).

In this study, there was a significant decrease in total antioxidant levels in Patients group compared with a control group ( $P<0.001$ ). Low dietary intakes of antioxidant vitamins or reduced synthesis of non-dietary antioxidants such as albumin, bilirubin, glutathione peroxidase cholesterol and uric acid are likely to result in an oxidant-antioxidant imbalance that exacerbates inflammation and tissue damage [13]. This result is in agreement with results of [14]. There was a significant decrease in superoxide dismutase levels in Patients group compared with the control group ( $P<0.001$ ). Superoxide dismutase decreases levels of oxidative stress and improves endothelial dysfunction and diminishes the development of atherosclerosis [15]. The results of this study show that reduce in copper/zinc SOD (Cu/ZnSOD) could associate with increased vascular oxidative stress, which agree with the results of [17]. There was a significant decrease in Catalase levels in Patients group compared with the control group ( $P<0.001$ ). To balance basal ROS production, the cellular redox milieu is maintained by a number of key antioxidant enzymes that limit intracellular and extracellular accumulation of deleterious reactive oxygen metabolites. These antioxidant defense enzymes include the superoxide dismutases, catalase, which reduces hydrogen peroxide to water and others, and the result is in agreement with results of [17] (Table 3-5).

## Conclusion

Antioxidants like SOD and CAT have shown promise as possible therapy for prevention and treatment of Ischemic heart diseases.

## Acknowledgement

The authors would like to express thanks to Dr.Ban Jaber for here kind help in statistical analysis of this article.

## References

1. Urbaniak A, Molski M, Szeląg M: **Quantum-chemical calculations of the antioxidant properties of trans-p-coumaric acid and trans-sinapinic acid.** *Comput. Methods Sci. Technol* 2012, **18**:117-128.
2. Jozanov-Stankov O, Đurić J, Dobutović B, Isenović ER: **Determination of total antioxidant status (TAS) as a biochemical parameter in control of workers' health.** *Archives of Biological Sciences* 2009, **61**:375-382.
3. Hamid A, Aiyelaagbe O, Usman L, Ameen O, Lawal A: **Antioxidants: Its medicinal and pharmacological applications.** *Afr J Pure Appl Chem* 2010, **4**:142-151.
4. Szeląg M, Mikulski D, Molski M: **Quantum-chemical investigation of the structure and the antioxidant properties of  $\alpha$ -lipoic acid and its metabolites.** *Journal of molecular modeling* 2012, **18**:2907-2916.
5. Mikulski D, Szeląg M, Molski M, Górniak R: **Quantum-chemical study on the antioxidation mechanisms of trans-resveratrol reactions with free radicals in the gas phase, water and ethanol environment.** *Journal of Molecular Structure: THEOCHEM* 2010, **951**:37-48.
6. Tietz NW: *Clinical guide to laboratory tests*: WB Saunders Co; 1995.
7. Herspring KF, Ferreira LF, Copp SW, Snyder BS, Poole DC, Musch TI: **Effects of antioxidants on contracting spinotrapezius muscle microvascular oxygenation and blood flow in aged rats.** *Journal of Applied Physiology* 2008, **105**:1889-1896.
8. Hirai DM, Copp SW, Schwagerl PJ, Haub MD, Poole DC, Musch TI: **Acute antioxidant supplementation and skeletal muscle vascular conductance in aged rats: role of exercise and fiber type.** *American Journal of Physiology-Heart and Circulatory Physiology* 2011, **300**:H1536-H1544.
9. Cohen G, Dembiec D, Marcus J: **Measurement of catalase activity in tissue extracts.** *Analytical biochemistry* 1970, **34**:30-38.
10. Djoussé L, Rothman KJ, Cupples LA, Levy D, Ellison RC: **Serum albumin and risk of myocardial infarction and all-cause mortality in the Framingham Offspring Study.** *Circulation* 2002, **106**:2919-2924.
11. Meisinger C, Heier M, Landgraf R, Happich M, Wichmann H-E, Piehlmeier W: **Albuminuria, cardiovascular risk factors and disease management in subjects with type 2 diabetes: a cross sectional study.** *BMC health services research* 2008, **8**:226.
12. Ogawa T, Ishida H, Akamatsu M, Matsuda N, Fujiu A, Ito K, Ando Y, Nitta K: **Progression of aortic arch calcification and all-cause and cardiovascular mortality in chronic hemodialysis patients.** *International urology and nephrology* 2010, **42**:187-194.
13. Fukai T, Folz RJ, Landmesser U, Harrison DG: **Extracellular superoxide dismutase and cardiovascular disease.** *Cardiovascular research* 2002, **55**:239-249.
14. Franzini L, Ardigo D, Valtuena S, Pellegrini N, Del Rio D, Bianchi M, Scazzina F, Piatti P, Brighenti F, Zavaroni I: **Food selection based on high total antioxidant capacity improves endothelial function in a low cardiovascular risk population.** *Nutrition, Metabolism and Cardiovascular Diseases* 2012, **22**:50-57.
15. Glorieux C, Dejeans N, Sid B, Beck R, Calderon PB, Verrax J: **Catalase overexpression in mammary cancer cells leads to a less aggressive phenotype and an altered response to chemotherapy.** *Biochemical pharmacology* 2011, **82**:1384-1390.
16. Li H, Horke S, Förstermann U: **Oxidative stress in vascular disease and its pharmacological prevention.** *Trends in pharmacological sciences* 2013, **34**:313-319.

17. Loenn ME, Dennis JM, Stocker R: **Actions of “antioxidants” in the protection against**

**atherosclerosis.** *Free Radical Biology and Medicine* 2012, 53:863-884.

Note: VRI Press, Vedic Research Inc. is not responsible for any data in the present article including, but not limited to, writeup, figures, tables. If you have any questions, directly contact authors.

Visit us @ [www.vedicjournals.com](http://www.vedicjournals.com) : DOI: <http://dx.doi.org/10.14259/bmc.v3i1.157>

Copyright © 2013-2014 VRI Press. All rights reserved.

## Authors Column



**Prof. Dr. Mufeed Jalil Ewadh**, Ph.D (Clinical Biochemistry), University of Wales, UK-1989, M.Sc (Chemistry) Honors, 1980-Iraq, B.Sc. (Chemistry) Honors, 1978-Iraq, Specialization, Clinical Enzymology, Protein Electrophoresis, Natural product Extraction for diseases treatments, Endothelial cell toxicity, Academic Staff of Medicine College, Babylon University, More than 110 papers. Recent activity, HbA1c training certificate, Marburg University, Germany



**Professor ( Dr ) monem Makki Alshok**

MBChB , MD , CABM , FRCP, Professor in clinical medicine college of medicine, Babylon University  
Consultant physician in Merjan teaching hospital Babylon, Chief of department of medicine college of medicine Babylon University till end of 2014 & now Supervisor of higher medical education in the college. Member of Iraqi medical board for medical specialties, and Supervisor of Postgraduate Babylon training center for medical specialties. Member of I.S.G.H. in Iraq & Member of Iraqi Diabetes association and Cardiology & chest diseases association and also member of ESGE (European Society of GI endoscopies).