

VRI Press Bioinformatics & Proteomics

eISSN 2330-0272

JOURNAL HOME PAGE AT WWW.VEDICJOURNALS.COM

Research Article

DOI: http://dx.doi.org/10.14259/bp.v3i1.125

Novel Antiviral Activity of Anthocyanins: An In Silico Approach

SHAILIMA DV RAMPOGU

Department of Biochemistry, Cachet Labs, Yousufguda, Hyderabad, Telangana, India

Article Info: Received: May 5th, 2014; Accepted: May 25th, 2014

ABSTRACT

HIV-1 protease [E.C.3.4.23.16] is an important enzyme involved in life cycle of HIV—a retrovirus causing AIDS. It cleaves the nascent polyprotein to generate the mature proteins of HIV virions, and its absence renders HIV uninfectious. Thus the HIV protease seems to be an interesting drug target for HIV. Utilising this property, this investigation aims to design a drug from the naturally available sources such as anthocyanins. Anthocyanins belong to flavonoid group of phytochemicals naturally present in tea, honey, wines, fruits, nuts, etc. . In this study, 10 anthocyanin compounds were designed and docked with HIV protease. In docking results, cyanidin, showing maximum dock score of 41.098, was found to be the best drug ligand.

Keywords: HIV, HIV Protease, Anthocyanins, Flavonoid groups, Phytochemicals.

INTRODUCTION

The term anthocyanin is derived from Greek word (flower) and kyanose (blue) to describe the blue pigment of Centaurea cyanus[1-2]. Anthocyanins are plant pigments soluble in water and primarily present in fruits and vegetables having bright colours [3]. They belong to the phenolic compounds and around 500 structures have been reported by year 2000 [4]. Flavonoids are important C₁₅ aromatic compounds of plant origin [5]. Chemically anthocyanins are glycoslyated, polyhydroxy or polymethoxy derivatives of 2-phenyl benzopyrlium containing two benzovl rings, separated by a heterocyclic ring [2]. These plant components are known to exhibit several medically important biological activities such as antioxidative and radical scavenging activity [6-10], improving sight [11-12] and anti-diabetic properties [13-14]. Besides these, they can also enhance insulin secretion [15-16] and can act as chemoprotective agents [17-18]. They also show antiviral [19] and antimicrobial activity [20] and cure common cold [21]. In summary, anthocyanins, as an important group of phenolic compounds, have wide medical applications beneficial to humans [22]. The most common six known anthocyanins are

*Corresponding Author

Shailima DV Rampogu

Department of Biochemistry, Cachet Labs, Yousufguda, Hyderabad, Telangana, India.

Email: shailima@cachetlabs.com

the pelargonidin, cyanidin, delphinidin, peonidin, petunidin and Malvidin. The daily intake of dietary flavonoids is estimated to be about 20-25 mg/day, while, especially in US it is between 180-215 mg/day [23].

The objective of the present experiment is to dock the anthocyanins with HIV protease (PDB ID: 1a8g)—a potential HIV drug target. This study focuses on the antiviral activity of the anthocyanins.

METHODS

The objective of the experiment is accomplished by preparing the ligand and the protein for the docking studies. Following the Protein-Ligand docking method, interactions were conducted and results were analysed that gave the flexibility of docking based on the structure.

A. Ligand Preparation

Anthocyanins were used as ligands for the docking studies and their structures were drawn using Chemsketch (ACD LABS 12.0). The duplicates were removed and bonds were added. The CHARMm force field was applied to minimise the energy.

B. Protein Preparation

The structure of the protein (ID: 1a8g) was imported from the



Rampogu, 2015

Protein Data Bank of high resolution to the Discovery Studio 2.5. The hetero-atoms and water molecules were removed from the structure. The chemistry of the missing hydrogens was corrected. By using alternate conformations and valence monitor options, crystallographic disorders and unfilled valence atoms were corrected.

C. Docking

The Protein-Ligand method was used for docking, which is based on molecular modelling. This method predicts the position and orientation of a small molecule called ligand when it binds to a receptor molecule, a protein or enzyme [24]. The docking was done on the Discovery Studio 2.5.




Figure 1: Active site pocket - Identification of Active site pocket by using Eraser algorithm in Discovery Studio.

RESULTS AND DISCUSSION

Ligand Preparation


The ligands for the present study, anthocyanins were drawn in Chemsketch (ACD LABS 12.0). Ten ligands were drawn [25] and their 3D structures and sterioisomers were generated.

Protein Preparation

The protein (ID: 1a8g) was imported from the Protein Data Bank. The energy minimisation of the protein was done using the conjugate gradient method and steepest descent method till the convergence gradient was satisfied. The active site pockets were identified by using the Eraser Algorithm and a sphere was created around the active site.

Protein-Ligand Docking

The results of the protein ligand docking studies were analysed, which showed cyanidin and two of its sterioisomers to be the best compound exhibiting antiviral properties with the highest dock score of 41.098, 41.094 and 40.928, respectively.



Amino acids involved in the interactions Gly27, Asp29, Cys95, Asp30

Amino acids involved in the interactions Asn98, Thr96, Glu2

Figure 2: H- bond interactions of Cyanidin with active site pocket residue of HIV protease.



Table 1: Docking Results - Table showing the dock results of Protein-Ligand Interaction

Name	Index	LigScore1_Dreiding	LigScore2_Dreiding	-PLP1	-PLP2	Jain	-PMF	DOCK_SCORE T
1 cyanidin	11	1.72	4.6	72.62	53.26	-0.35	15.39	41.098
2 cyanidin	12	1.74	4.56	71.61	52.58	-0.21	14.4	41.094
3 cyanidin	13	1.74	4.56	71.61	52.58	-0.21	14.4	41.094
4 cyanidin	14	1.69	4.45	72.14	53.75	-0.17	13.33	40.928
5 cyanidin	15	1.59	4.58	64.96	54.62	-0.32	7.56	40.692
6 cyanidin	16	1.59	4.58	64.96	54.62	-0.32	7.56	40.692
7 aurantinidin	1	1.65	4.69	63.94	55.12	-0.39	9.55	40.608
8 aurantinidin	2	1.74	4.6	69.94	54.23	-0.45	12.57	40.583
9 cyanidin	17	1.45	4.62	62.54	49.49	-0.73	12.82	40.569
10 cyanidin	18	1.62	4.58	63.98	54.57	-0.26	9.6	40.562
11 cyanidin	19	1.6	4.63	60.61	52.56	-0.51	9.65	40.506
12 cyanidin	20	1.6	4.63	60.61	52.56	-0.51	9.65	40.506
13 luteolinidin	31	1.52	4.69	63.4	55.78	-0.08	10.33	40.48
14 luteolinidin	32	1.52	4.69	63.4	55.78	-0.08	10.33	40.48
15 aurantinidin	3	1.62	4.72	62.34	52.74	-0.52	9.89	40.447
16 aurantinidin	4	1.61	4.7	62.14	54.24	-0.42	8.74	40.411
17 aurantinidin	5	1.61	4.7	62.14	54.24	-0.42	8.74	40.411
18 aurantinidin	6	1.61	4.7	62.14	54.24	-0.42	8.74	40.411
19 aurantinidin	7	1.61	4.7	62.14	54.24	-0.42	8.74	40.411
20 luteolinidin	33	1.52	4.71	63.76	55.27	0	9.89	40.311
21 luteolinidin	34	1.52	4.71	63.76	55.27	0	9.89	40.311
22 luteolinidin	35	1.52	4.71	63.76	55.27	0	9.89	40,311
23 luteolinidin	36	1.52	4.71	63.76	55.27	0	9.89	40.311
24 luteolinidin	37	1.53	4.64	63.08	54.55	-0.02	11.27	40.258
25 aurantinidin	8	1.56	4.64	62.75	53.75	-0.66	11.01	40.218
26 aurantinidin	9	1.54	4.65	62.99	53.05	-0.73	10.9	40.151
27 aurantinidin	10	1.54	4.65	62.99	53.05	-0.73	10.9	40.151
28 luteolinidin	38	1.53	4.61	60.81	53.34	-0.01	10.14	40.064
29 luteolinidin	39	1,53	4.61	60.81	53.34	-0.01	10.14	40.064
30 luteolinidin	40	1.46	4.66	65.8	56.47	-0.21	10.1	40.049
31 pelargonidin	41	1.67	4.56	62.66	54.16	-0.28	8.54	39.934




Figure :3 Showing the electrostatic surface area.



Conclusions

Anthocyanins the naturally available plant components are known to exhibit several medical properties. The present article successfully accomplishes the antiviral activity of the anthocyanins. Amongst the available anthocyanins, cyanidin proves to be the most lucrative drug against HIV.

REFERENCES

- Dilip Ghosh and Tetsuya Konishi: Anthocyanins and anthocyanin-rich extracts: role in diabetes and eye function. Asia Pac J Clin Nutr 2007; 16: 200-208.
- Jaclyn Shipp and El-Sayed and M. Abdel-Aal: Food Applications & Physiological Effects of Anthocyanins as Functional Food Ingredients. The open Food Science 2010, 4:7-22.
- Li-Shu Wang and Gary D. Stoner: Anthocyanins and their role in cancer prevention. Cancer Lett. 2008, 269: 281-290. doi: 10. 1016/j. canlet. 2008.05.020.
- Pietta PG: Flavonoids as antioxidants. J Nat Product 2000, 63: 1035-1042. DOI: 10.1021/np9904509
- Mary Ann Lila: Anthocyanins and Human Health: An In vitro investigative Approach. J Biomed Biotechnol 2004, 5: 306– 313. doi: 10.1155/S111072430440401X
- Abdel-Aal E S, Abou-Arab AA, Gamel TH, Hucl P, Young JC, Rabalski I: Fractionation of blue wheat anthocyanin compounds and their contribution to antioxidant properties. J Agric Food Chem 2008, 56: 11171-11177. DOI: 10.1021/jf802168c
- 7. Astadi IR, Astuti M, Santoso U, Nugraheni PS: Invitro antioxidant activity of anthocyanins of black soybean seed coat in human low density lipoprotein (LDL). Food Chem 2009, 112: 659-663. DOI: 10.1016/j.foodchem.2008.06.034
- Bellido GG; Beta T: Anthocyanin composition and oxygen radical scavenging capacity (ORAC) of milled & pearled purple, black and common barley. J. Agric Food chem. 2009, 57: 1022-1028. DOI: 10.1021/jf802846x
- 9. Kahkonen MP, Heinonen M: Antioxidant activity of anthocyanins and their aglycones, J. Agric Food Chem. 2003, 51: 628-633. DOI: 10.1021/jf025551i
- Sanchez-Moreno C, Satue-Gracia MT, Frankel EN: Antioxidant activity of selected sapnish wines in corn oil emulsion. J Agric Food chem, 2008, 48: 5581-5587. DOI: 10.1021/ jf000664s
- 11. Ghosh D., and Konishi T: **Anthocyanins and anthocyanin-rich** extracts: role in diabetes and eye function. *Asia Pac J Clin Nutr* 2007, **16**: 200-208. PMID:17468073
- 12. Kalt W, Blumberg JB, Mcdonald JE *et al*: **Identification of anthocyanins in the liver, eye and brain of blueberry fed pigs.** *J Agric Food Chem* 2008, **56**: 705-712. doi: 10.1021/jf071998l

- 13. De Furia J, Bennett G, Strissed KJ, et al: Dietary blueberry attenuates whole-body insulin resistance in high-fat fed mice by reducing adipocyte death and its inflammatory sequelae, *J. Nutr.* 2009, 139: 1-7. doi: 10.3945/jn. 109.105155
- 14. Rosenblat M, Hayek T, Aviram M: Anti-oxidative effects of pomegranate juice(PJ) consumption by diabetic patients on serum and on macrophages. Antherosclerosis 2006, 187: 363-371. doi:10.1016/j.atherosclerosis.2005.09.006
- 15. Jayaprakasam B, Vareed Sk, Olson LK, Nair MG: Insulin secretion by bioactive anthocyanins and anthocyanidins present in fruits. *J Agric Food Chem* 2005, 53: 28-31. DOI: 10.1021/jf049018+
- 16. Zhang Y, Jayaprakasam B, Seeram NP, Olson LK, Dewitt D, Nair MG: Insulin secretion and cyclooxygenase enzyme inhibition by cabernet sauvignon grape skin compounds. J Agric Food Chem 2004, 52: 228-233. DOI: 10.1021/jf034616u
- 17. Colditz GA, Branch LG, Lipnick RJ et al: Increased green and yellow vegetable intake and lowered cancer deaths in an elderly population. Am J Clin Nutr 1985, 41: 32-36. PMID: 3966422
- 18. Liu Z, Schwimer J, Liu D, Greenway FL, Anthony CT, Woltering EA: Black raspberry extract and fractions contain angiogenesis inhibitors. *J Agric Food Chem* 2005, 53: 3909-3915. DOI: 10.1021/jf048585u
- Cheng HY, Lin CC, Lin TC: Antiviral properties of prodelphinidin B-2 3'-O-gallate from green tea leaf. Antivir Chem Chemother. 2002, 13: 223-9 PMID:12495210.
- Cisowska A, Wojnicz D, Hendrich AB: Anthocyanins as antimicrobial agents of natural plant origin. Nat Prod Commun. 2011, 6:149-56.PMID:21366068.
- Konczak I, Zhang W: Anthocyanins- more than nature's colours.
 J Biomed Biotechol 2004, 5: 239-240.doi: 10.1155/ S1110724304407013
- 22. Simona Oancea, Letitia Oprean Anthocyanins, from biosynthesis in plants to human health benefits, Acta universitatis Cibiniensis Series E: Food Technology vol. XV (2011), no. 1; http://saiapm.ulbsibiu.ro/rom/cercetare/ACTA_E/AUCFT_2011_I_3_16.pdf; Last accessed: 30-April-2014.
- Hertog MG, Hollman PC, Katan MB, Kromhout D: Intake of potentially anticarcinogenic flavonoids and their determinants in adults in the Netherlands. Nutr Cancer 1993, 20: 21-29. Doi: 10.1080/01635589309514267
- 24. Alejandra Hernández-Santoyo, Aldo Yair Tenorio-Barajas, Victor Altuzar, Héctor Vivanco- Cid and Claudia Mendoza-Barrera (2013). Protein-Protein and Protein- Ligand Docking, Protein Engineering Technology and Application, Dr. Tomohisa Ogawa (Ed.), ISBN: 978-953-51-1138-2, InTech, DOI: 10.5772/56376.
- 25. Vardhini SRD: In silico evaluation for the potential naturally available drugs for the brast cancer. *Journal of Receptors and Signaling Transduction* 2014, 3: 174-179.

<u>Note:</u> VRI Press, Vedic Research Inc. is not responsible for any data in the present article including, but not limited to, writeup, figures, tables. If you have any questions, directly contact authors.

Visit us @ www.vedicjournals.com : DOI: http://dx.doi.org/10.14259/bp.v3i1.125

Copyright © 2013-2015 VRI Press, USA. All rights reserved.

